

CHEST[®]

Official publication of the American College of Chest Physicians



Pulmonary Complications of Novel Antineoplastic Agents for Solid Tumors

Bobbak Vahid and Paul E. Marik

Chest 2008;133:528-538
DOI 10.1378/chest.07-0851

The online version of this article, along with updated information and services can be found online on the World Wide Web at:
<http://chestjournal.org/cgi/content/abstract/133/2/528>

CHEST is the official journal of the American College of Chest Physicians. It has been published monthly since 1935. Copyright 2007 by the American College of Chest Physicians, 3300 Dundee Road, Northbrook IL 60062. All rights reserved. No part of this article or PDF may be reproduced or distributed without the prior written permission of the copyright holder (<http://www.chestjournal.org/misc/reprints.shtml>). ISSN: 0012-3692.

A M E R I C A N C O L L E G E O F



P H Y S I C I A N S[®]



Pulmonary Complications of Novel Antineoplastic Agents for Solid Tumors*

Bobbak Vahid, MD; and Paul E. Marik, MD, FCCP

Antineoplastic agent-induced pulmonary toxicity is an important cause of respiratory failure. Although the incidence of antineoplastic agent-induced pulmonary toxicity seems to be low, more cases can be expected, with increasing numbers of patients receiving the new generations of antineoplastic agents. Antineoplastic agents have previously been associated with bronchospasm, hypersensitivity reactions, venous thromboembolism, and pulmonary hemorrhage. Physicians should be aware of the clinical and radiographic presentations of the pulmonary toxicities associated with the newer antineoplastic agents. The approach to diagnosis, risk factors, and possible mechanisms of antineoplastic agent-induced pulmonary toxicity are discussed in this article. (*CHEST* 2008; 133:528–538)

Key words: chemotherapy; pneumonitis; pulmonary toxicity; respiratory failure

Abbreviations: CML = chronic myelogenous leukemia; DVT = deep venous thrombosis; EGFR = epidermal growth factor receptor; HER = human epidermal growth factor receptor; KL-6 = Krebs von den Lunge-6; mTOR = mammalian target of rapamycin; NSCLC = non-small cell lung cancer

Antineoplastic agent-induced pulmonary toxicity is an important cause of respiratory failure. New antineoplastic agents and regimens are constantly being added to the list of available treatments for cancer patients. These novel antineoplastic agents either have new mechanisms of action such as tyrosine kinase inhibitors or old agents with new indications like thalidomide. Since more patients are being treated with these agents, associated acute respiratory failure is more commonly being recognized. Critical care physicians should be aware of the clinical and radiographic presentations of antineoplastic agent-induced pulmonary toxicities. Unfortunately, the diagnosis of antineoplastic agent-induced pulmonary toxicity is complicated. Antineoplastic

agent-induced pulmonary toxicity is a diagnosis of exclusion, and other causes of respiratory failure including pneumonia, cardiogenic pulmonary edema, and diffuse alveolar hemorrhage should be excluded. These conditions are not easily differentiated based on clinical presentation and radiographic findings. Furthermore, as patients usually receive multiple antineoplastic agents, it is usually difficult to identify the culprit agent. Open-lung biopsy may be necessary in selected cases to exclude the alternative diagnoses. Regardless of these difficulties, antineoplastic agent-induced pneumonitis and respiratory failure should be considered in patients receiving chemotherapeutic agents. The cessation of the implicated causative agent and treatment with systemic corticosteroids may result in rapid improvement.^{1,2}

CLINICAL MANIFESTATIONS AND DIAGNOSIS

Several clinical syndromes have been described in patients with presumed antineoplastic agent-induced lung toxicity (Table 1). The definition of these clinical syndromes may be confusing due to the different criteria used in the literature. Most clinical trials do not report the details of pulmonary toxicity. Authors describe the pulmonary toxicities based

*From the Department of Pulmonary and Critical Care Medicine, Thomas Jefferson University Hospital, Philadelphia, PA. The authors have reported to the ACCP that no significant conflicts of interest exist with any companies/organizations whose products or services may be discussed in this article. Manuscript received April 5, 2007; revision accepted May 18, 2007.

Reproduction of this article is prohibited without written permission from the American College of Chest Physicians (www.chestjournal.org/misc/reprints.shtml).

Correspondence to: Bobbak Vahid, MD, 834 Walnut St, Suite 650, Philadelphia, PA 19107; e-mail: bobbak.vahid@mail.tju.edu
DOI: 10.1378/chest.07-0851

Table 1—Definitions of Clinical Syndromes*

Syndromes	Description
Bronchospasm	Evidence of airflow obstruction (<i>eg</i> , wheezing and prolonged exhale)
Hypersensitivity reaction	Bronchospasm plus other hypersensitivity-related symptoms: angioedema; rash; urticaria; hypotension; arthralgia; nausea; vomiting; hypotension; or hypertension
Infusion reaction	Hypersensitivity reaction during infusion or shortly (within minutes) thereafter
Pneumonitis	Clinical and radiographic manifestation compatible with interstitial pneumonitis
Noncardiogenic pulmonary edema	Pulmonary edema not associated with heart failure or increased left atrial pressures
Capillary leak syndrome	Noncardiogenic pulmonary edema associated with diffuse edema and intravascular hypovolemia
Acute lung injury	1. Noncardiogenic pulmonary associated with evidence of acute inflammation like: fever; and elevated neutrophils in BAL fluid 2. Diffuse bilateral pulmonary infiltrates, no evidence of elevated left atrial pressure, and PO_2/FIO_2 ratio < 300
ARDS	Diffuse bilateral pulmonary infiltrates, no evidence of elevated left atrial pressure, and PO_2/FIO_2 ratio < 200
Eosinophilic pneumonia	Pulmonary infiltrates, hypoxemia, and BAL fluid eosinophilia

* FIO_2 = fraction of inspired oxygen.

either on clinical criteria (*eg*, acute lung injury, ARDS, noncardiogenic pulmonary edema, or pneumonitis) or pathologic findings (*eg*, diffuse alveolar damage, organizing pneumonia, nonspecific pneumonitis, or neutrophilic alveolitis). In this review, we will use the definitions outlined in Table 1.

The clinical manifestations of drug-induced pneumonitis are nonspecific and include cough, fever, dyspnea, and hypoxemia. The pulmonary involvement may be rapidly progressive, resulting in respiratory failure and ARDS. The timing of clinical manifestations is unpredictable; they may present during the first cycle of treatment or following subsequent cycles.^{1,2} Concurrent treatment with corticosteroids may not prevent the development of drug-induced pneumonitis.³⁻⁷ Elevated WBC count in peripheral blood, elevated erythrocyte sedimentation rate, and elevated C-reactive protein levels are also common nonspecific laboratory findings. Chest imaging may show diffuse or patchy, unilateral or bilateral, ground-glass opacities or consolidations. Reduced lung volumes and diffusion capacity are commonly reported. BAL fluid cell counts are usually elevated with neutrophilia, lymphocytosis, or, rarely, eosinophilia.^{1,2}

Elevated serum Krebs von den Lunge-6 (KL-6) levels have been reported in patients with drug-

induced pneumonitis. KL-6 is a mucin-like high-molecular-weight glycoprotein that is expressed by type II alveolar pneumocytes. Elevated levels of KL-6 (*ie*, ≥ 500 U/mL) can also be seen in patients with idiopathic interstitial pneumonitis, pneumonitis related to collagen vascular disease, and hypersensitivity pneumonitis.⁸ Furthermore, only 53.3% of patients with drug-induced pneumonitis have elevated serum KL-6 levels.⁹ However, KL-6 levels do not increase in patients with pneumonia, pulmonary aspergillosis, asthma, bronchiectasis, emphysema, eosinophilic pneumonia, or organizing pneumonia.⁸ Other causes of elevated KL-6 levels include hypersensitivity pneumonitis, radiation pneumonitis, viral pneumonia, pneumocystis pneumonia, sarcoidosis, tubulointerstitial nephritis uveitis syndrome, liver disease (*eg*, hepatitis C, cirrhosis, or hepatocellular carcinoma), and malignancy (*eg*, breast cancer or malignant thymoma).⁸⁻¹⁵

One unique presentation of antineoplastic agent-induced pneumonitis is so-called *radiation recall pneumonitis*. Radiation recall pneumonitis is seen in patients who have received previous radiation therapy to the chest. Shortly after the initiation of therapy with the antineoplastic agent, the fever, cough, dyspnea, and hypoxemia develop in the patient. The chest imaging shows pulmonary infiltrates in exactly the same field as in previous radiation therapy. Subsequent progression to diffuse bilateral pneumonitis can be expected in severe cases. The exact mechanism of radiation recall pneumonitis is not known. Irradiation may cause subclinical injury to the lung parenchyma, which may have an additive effect in precipitating lung injury when another pulmonary insult is encountered at a later date. Other possible mechanisms include injury to type II pneumocytes in the radiotherapy field, which reduces the ability of the lung to repair itself and drug hypersensitivity reactions. Other manifestations of radiation recall are dermatitis, mucositis, and myositis. The antineoplastic agents associated with radiation recall pneumonitis are adriamycin, carmustine, doxorubicin, etoposide, gefitinib, gemcitabine, paclitaxel, and trastuzumab.¹⁶⁻²²

The differential diagnosis of antineoplastic agent-induced pneumonitis is extensive. Infection is a very common cause of pulmonary infiltrates and respiratory failure in cancer patients. Appropriate cultures and serology can be helpful to differentiate pneumonitis from infectious pneumonia. Bronchoscopy with BAL is very useful to exclude an infectious process. Bronchoscopy with BAL is also important to exclude alveolar hemorrhage. In patients with alveolar hemorrhage, the BAL fluid return is hemorrhagic and contains hemosiderin-laden macrophages. The presence of malignant cells in BAL fluid suggests the

lymphangitic spread of the cancer. ARDS secondary to aspiration, sepsis, and pancreatitis may be suspected based on history, physical examination results, and other laboratory workup findings. Cardiogenic pulmonary edema is suggested if the echocardiogram shows left ventricular dysfunction and serum B-type natriuretic peptide is elevated. Bilateral transudative pleural effusions and rapid improvement in pulmonary status after diuresis are typically observed in patients with cardiogenic pulmonary edema.^{1,2}

Transbronchial or open-lung biopsy can be helpful in demonstrating the presence of pneumonitis and excluding alternative diagnoses, like lymphangitic carcinomatosis, vasculitis, and pneumonia. Nonspecific pneumonitis, organizing pneumonia, eosinophilic pneumonia, pulmonary fibrosis, and diffuse alveolar damage are commonly seen on lung biopsy specimens. These pathologic patterns are nonspecific and should not be considered diagnostic of drug-induced lung disease. The diagnosis of chemotherapy-induced pneumonitis can be made when pneumonitis develops shortly after the initiation of treatment (*ie*, hours to weeks), lack of an alternative explanation for respiratory failure, and the resolution of pneumonitis after corticosteroid treatment and withdrawal of the presumed agent.

PATHOGENESIS AND RISK FACTORS

The pathogenesis of antineoplastic agent-induced lung injury is poorly understood. Several mechanisms have been suggested. Direct injury to pneumocytes (chemical alveolitis) or the alveolar capillary endothelium and the subsequent release of cytokines and recruitment of inflammatory cells may be responsible, along with some of the cytotoxic medications. The systemic release of cytokines by chemotherapeutic agents (*eg*, gemcitabine) may also result in capillary leak and pulmonary edema. Positive lymphocyte stimulation test results and an elevated CD4/CD8 cell ratio suggest cell-mediated lung injury due to the activation of lymphocytes, and alveolar macrophages may also play a role. Free oxygen radicals may also be involved especially with mitomycin-C pulmonary toxicity. Epidermal growth factor receptors (EGFRs) are expressed on type II pneumocytes and are involved in alveolar wall repair. EGFR tyrosine kinase inhibitors, by impairing the alveolar repair mechanisms, may potentiate the effect of lung injury due to other causes, including sepsis, radiation, and other medications.^{23–26}

Combination chemotherapy may have an additive effect with a higher frequency of pulmonary toxicity. Preexisting pulmonary disease such as idiopathic

pulmonary fibrosis, COPD, radiation therapy, extensive pulmonary metastatic disease, and poor functional status have also been associated with increased pulmonary toxicity. A high inspired oxygen concentration may increase the incidence or severity of mitomycin-C-induced pneumonitis.^{23,24}

ANTINEOPLASTIC AGENTS

Table 2 summarizes the pulmonary complications associated with the newer antineoplastic agents used in the treatment of solid tumors.^{27–112} Each agent is discussed separately below.

ALKYLATING AGENTS

Chlorozotocin

Chlorozotocin is an alkylating agent with activity against advanced islet-cell carcinoma. Two reported cases^{28,29} of mild subacute interstitial pneumonitis have been reported with this agent. Mild dyspnea, dry cough, and pulmonary infiltrates developed in one patient after the third dose of chlorozotocin. Pulmonary function test results showed reduced diffusion capacity and lung volumes. Acute pneumonitis developed in the second patient after receiving chlorozotocin and mitomycin-C. Pneumonitis resolved with prednisone therapy with no residual pulmonary fibrosis.^{28,29}

Ifosfamide

Ifosfamide is an alkylating agent that is active against a variety of solid tumors including breast cancer, lung cancer, ovarian cancer, and sarcomas. Ifosfamide pulmonary toxicity has been noted in combination with other antineoplastic therapies. Fatal interstitial pneumonitis has been reported with ifosfamide therapy for soft-tissue sarcoma.³⁰ In a phase II trial³¹ of docetaxel and ifosfamide, interstitial pneumonitis developed after combination chemotherapy in three patients (6%) with non-small cell lung cancer (NSCLC). Two of three patients died due to respiratory failure.³¹ The ifosfamide metabolite, 4-thioifosfamide, is known to react with glutathione and deplete the RBC antioxidant reserve. This reaction may rarely result in methemoglobinemia.³²

Oxaliplatin

Oxaliplatin is a new cytotoxic agent that is mainly used in the treatment of colorectal cancer combined with fluorouracil and leucovorin. Interstitial pneumonitis with fibrosis has been reported after 3 to 6

Table 2—Pulmonary Complications of Chemotherapeutic Agents

Chemotherapeutic Agents	Pulmonary Complications
Bevacizumab	Pulmonary hemorrhage and hemoptysis Increased risk of DVT and pulmonary embolism
Chlorozotocin	Interstitial pneumonitis
Doxorubicin	Organizing pneumonia Infusion reaction
Erlotinib	Acute pneumonitis ARDS
Etoposide	Acute pneumonitis Diffuse alveolar damage Hypersensitivity reaction/bronchospasm
Everolimus	Acute pneumonitis
Gefitinib	Acute pneumonitis Diffuse alveolar damage Diffuse alveolar hemorrhage Pulmonary fibrosis
Gemcitabine	Infusion-related reaction Capillary leak syndrome/pulmonary edema ARDS Diffuse alveolar damage Diffuse alveolar hemorrhage Bronchospasm Pleural effusion
Ifosfamide	Interstitial pneumonitis Methemoglobinemia
Imatinib	Acute pneumonitis Fluid retention and pulmonary edema Pleural effusion
Irinotecan	Moderate-to-severe pneumonitis Severe hypoxemia and respiratory failure
Oxaliplatin	Pulmonary fibrosis Respiratory failure Eosinophilic pneumonia Infusion-related reactions
Matuzumab	Bronchospasm
Mitozantrone	Acute pneumonitis
Piritrexim	Acute pneumonitis
Taxanes	Acute pneumonitis Infusion-related reactions Pleural effusion
Temozolomide	Acute pneumonitis
Temsirolimus	Acute pneumonitis
Thalidomide	Pulmonary embolism Organizing pneumonia Acute pneumonitis Pleural effusion
Teniposide	Hypersensitivity reaction/bronchospasm
Topotecan	Bronchiolitis and organizing pneumonia
Trastuzumab	Acute lung injury Acute pneumonitis Organizing pneumonia Infusion-related reaction Bronchospasm

months of therapy. The patients present with slowly progressive cough and dyspnea for several months, but the disease can become accelerated. Deaths due to respiratory failure have been reported 10 to 20 days after presentation even with corticosteroid ther-

apy.^{33,34} Eosinophilic pneumonia after oxaliplatin therapy is a rare complication.³⁵

Oxaliplatin infusion-related reactions and severe anaphylactic reactions occur with a reported frequency of 1.3%. All reactions appear within 5 to 50 min after starting oxaliplatin infusion and usually last less than a day. A hypertensive crisis resulting in a change in mental status has been reported.³⁶

Temozolomide

Temozolomide is a new alkylating agent with preclinical activity against a variety of solid tumors and hematologic malignancies. It is mainly used in the treatment of anaplastic astrocytoma and metastatic melanoma. There are limited data from phase I and II trials^{37,38} regarding the safety and efficacy of temozolomide in the treatment of malignancies. In a phase II trial³⁷ of temozolomide for patients with recurrent or progressive brain metastases from a variety of solid tumors, pneumonitis developed in 4.8% of patients. The patients were treated with 150 to 200 mg/m²/d for 5 days. The treatment cycles were repeated every 4 weeks. In another phase II trial,³⁸ 46 elderly patients with acute myeloid leukemia were treated with oral temozolomide, 200 mg/m²/d for 5 days. One patient died of acute interstitial pneumonitis.^{37,38}

ANTIBIOTICS

Doxorubicin

Doxorubicin is a cytotoxic antibiotic that inhibits topoisomerase II. It has activity against a variety of solid tumors (*ie*, cancers of the bladder, breast, stomach, lung, ovaries, and thyroid, soft-tissue sarcoma, and others). Lung toxicity is rare. Infusion reaction may be seen in 8% of patients during pegylated liposomal doxorubicin infusion. Dyspnea may develop in patients within 1 to 5 min after infusion, and the symptoms resolve within 5 to 15 min after stopping the infusion. *In vitro* studies³⁹ have shown that pegylated-liposomal doxorubicin stimulates neutrophil adhesion to human umbilical vein endothelial cells. Since transient relative neutropenia has been detected during pegylated-liposomal doxorubicin infusion, the adhesion and sequestration of neutrophils to the pulmonary circulation have been suggested as a potential mechanism for infusion-related acute dyspnea.³⁹ Several cases of doxorubicin-induced organizing pneumonia, mostly in patients with lymphoma, have been described in the literature.^{40,41}

Epirubicin

Epirubicin is primarily used against breast and ovarian cancer, gastric cancer, lung cancer, and

lymphomas. Epirubicin pulmonary toxicity has been noted in combination with other antineoplastic therapies. Although rare, severe pneumonitis can occur in patients receiving epirubicin within weeks after undergoing irradiation to chest. In one study,⁴² interstitial pneumonitis occurred in 9% of patients (4 of 44 patients) with breast cancer treated with 5-fluorouracil, epirubicin, and cyclophosphamide. It is important to note that patients were also treated with granulocyte-stimulating factor, and two patients received one dose of paclitaxel.⁴² Although these reports do not support the pulmonary toxicity of epirubicin *per se*, epirubicin may potentiate the pulmonary toxicity of other agents.

Mitoxantrone

Mitoxantrone is a topoisomerase II inhibitor that is mainly used in the treatment of metastatic breast cancer. Severe acute pneumonitis may be seen after mitoxantrone therapy. The lung pathology demonstrates organizing pneumonia and a hypersensitivity pneumonitis-like pattern. Other chemotherapy agents (*ie*, bleomycin, vincristine, cyclophosphamide, and chlorambucil) were also included in the chemotherapeutic regimen in most cases. Oral corticosteroids were reported to result in the rapid resolution of pneumonitis.^{43–45}

ANTIMETOBOLITES

Piritrexim

Piritrexim is an orally bioavailable, second-generation antifolate with activity against transitional cell carcinoma. In a phase II trial,⁴⁶ piritrexim was administered orally in a dose of 25 mg daily for 5 consecutive days per week for 3 consecutive weeks. In this study,⁴⁶ pulmonary toxicity (only 1 of 28 patients had grade 3 pulmonary toxicity) developed in 14% of patients (4 of 28 patients).⁴⁶ de Wit et al⁴⁷ described a case of diffuse interstitial pneumonitis and respiratory failure after treatment with piritrexim for transitional cell carcinoma of the renal pelvis. The authors concluded that pulmonary disease was probably induced by piritrexim. The respiratory failure resolved after drug discontinuation.⁴⁷

MONOCLONAL ANTIBODIES

Bevacizumab

Bevacizumab, a monoclonal antibody against endothelial growth factor, has been used to treat patients with a variety of cancers. There are several reported pulmonary toxicities associated with bevacizumab therapy.

Pulmonary hemorrhage and hemoptysis has been reported^{48–50} in 2.3% of patients with nonsquamous NSCLC. In these patients, pulmonary hemorrhage may lead to respiratory failure, and fatalities have been reported in 1.6% of patients treated with bevacizumab. Severe hemoptysis and pulmonary hemorrhage associated with bevacizumab therapy is more common in patients, with squamous cell carcinoma being reported in up to 31% of patients. Bevacizumab has also associated with increased risk of deep venous thrombosis (DVT) and pulmonary embolism.^{48–50}

Matuzumab

Matuzumab is a humanized Ig G1 that blocks the activation of EGFR. Matuzumab is active against EGFR-positive NSCLC. Bronchospasm related to matuzumab has been reported in 5% of patients. Premedication with corticosteroids is effective to prevent bronchospasm.⁵¹

Trastuzumab

Trastuzumab is a humanized monoclonal antibody that selectively binds to the human EGFR (HER)-2 protein. Trastuzumab is indicated for the treatment of metastatic breast cancers with overexpression of HER-2 protein. The incidence of trastuzumab-induced pneumonitis is 0.4 to 0.6%. Trastuzumab-induced pneumonitis may present with rapidly progressive pulmonary infiltrates and respiratory failure after receiving 1 dose of trastuzumab or after 6 weeks of therapy. The mortality of trastuzumab-induced pneumonitis is about 0.1%. Acute neutrophilic alveolitis and organizing pneumonia after trastuzumab treatment also have been reported. Infusion-related symptoms, including hypotension, angioedema, bronchospasm, dyspnea, fever, chills, and urticaria, has been reported to occur in about 15% of patients. Severe episodes of hypotension, bronchospasm, and hypoxemia leading to death are rare.^{52–56}

NUCLEOSIDE ANALOGS

Gemcitabine

Gemcitabine is a nucleoside analog with activity against a variety of solid tumors, especially NSCLC and pancreatic cancer. A variety of forms of pulmonary toxicity have been described with the use of gemcitabine. Dyspnea developing within hours of infusion has been reported to occur in about 10% of patients. Most patients improve with therapy with diuretics and corticosteroids. Bronchospasm develops in about 0.6% of patients. These infusion-related

reactions are usually mild and rarely have resulted in the discontinuation of treatment.²³ Gemcitabine-induced pneumonitis has been reported to occur in up to 13.8% of patients. An analysis^{57,58} of pooled data from a large database showed the incidence of gemcitabine-induced pulmonary toxicity to vary from 0.02 to 0.27%. The following three types of gemcitabine-induced pneumonitis have been described: (1) a capillary leak syndrome with pulmonary edema; (2) diffuse alveolar damage; and (3) alveolar hemorrhage. Previous lung disease, chest radiation, and concurrent treatment with other agents (*eg*, paclitaxel, docetaxel, ifosfamide, or granulocyte colony-stimulating factor) are possible risk factors. Restrictive physiology with a marked reduction in diffusion capacity has been reported. Although the mortality rate can be as high as 20%, a rapid response (within days) to prednisone, 60 mg daily, has been described in the literature.^{23,57–63}

PODOPHYLLOTOXINS

Etoposide

Etoposide is a topoisomerase II inhibitor. This agent is used primarily in the treatment of small cell lung cancer. The most common pulmonary toxicity is a hypersensitivity reaction that can present with symptoms of anaphylaxis, angioedema, chest discomfort, bronchospasm, and hypotension.⁶⁴ Etoposide-induced acute pneumonitis or acute lung injury, although uncommon, may occur. The pathology of etoposide-induced lung injury is diffuse alveolar damage, fibrin membrane formation, and alveolar wall edema. Fatal cases are rare. Etoposide is also known to increase the risk of radiation pneumonitis. Concurrent treatment with other pneumotoxic agents may increase the risk of pneumonitis.^{65,66} Zimmerman et al⁶⁷ reported that interstitial pneumonitis developed in 24% of 50 patients treated with etoposide, methotrexate, and cyclophosphamide for small cell anaplastic lung cancer. It has been shown that etoposide increases the intracellular levels of methotrexate. The authors postulated that the pulmonary toxicity was secondary to methotrexate pneumonitis, which was caused by excessive intracellular levels of methotrexate related to the concurrent use of etoposide.⁶⁷

Teniposide

Teniposide is used in the treatment of glioblastoma multiforme. Hypersensitivity reaction to teniposide occurs in 3.6 to 6.5% of patients. urticaria, hypotension or hypertension, dyspnea, bronchospasm, cyanosis, flushing, and vomiting within first 10

to 20 min of teniposide infusion develop in patients. In patients with leukemia, infusion reaction tends to occur after the completion of infusion. However, the timing of the hypersensitivity reaction is unpredictable and may occur during the first treatment cycle or during subsequent treatment cycles.⁶⁸

RAPAMYCIN ANALOGS

Temsirolimus

Temsirolimus is a rapamycin analog that is active against renal cell carcinoma, endometrial carcinoma, breast cancer, glioblastoma multiforme, and GI neuroendocrine tumors. Temsirolimus binds with immunophilin FK-506 binding protein-12 and forms a complex that inhibits the protein activity of mammalian target of rapamycin (mTOR). mTOR is a serine-threonine kinase that regulates cell growth, proliferation, and apoptosis. Interstitial pneumonitis is a non-dose-dependent complication of temsirolimus. Interstitial pneumonitis has been reported in 1 to 36% of patients treated with 25 to 250 mg/wk. The onset of pneumonitis usually takes place within 16 weeks (range, 2 to 16 weeks) after temsirolimus treatment. In one case series,^{69,70} 50% of patients were clinically asymptomatic, with drug-induced pneumonitis diagnosed by chest imaging. The chest CT scan may show ground-glass opacity or consolidation.^{69,70}

Everolimus

The pharmacologic effects of everolimus are also mediated through binding to FK-506 binding protein-12 and the inhibition of mTOR. Everolimus has been used as an immunosuppressive agent following organ transplantation, to treat severe psoriasis, and as an investigational antineoplastic agent (*eg*, for the treatment of sarcoma or renal cell cancer). Although clinical data in patients with malignancy are sparse, in one study⁷¹ using everolimus in heart transplant recipients, interstitial pneumonitis developed in 3.3% of patients 4 weeks after switching treatment to everolimus. All patients with pneumonitis required mechanical ventilation.⁷¹

TAXANES

Taxanes are mainly used in the treatment of breast, ovarian, and lung cancers. Paclitaxel and docetaxel are known to cause pneumonitis with estimated frequencies of 0.73 to 12% and 7 to 26%, respectively. Dyspnea, cough, hypoxemia, and pulmonary infiltrates usually develop 1 week to 3

months after treatment. Possible risk factors for pulmonary toxicity are weekly or biweekly therapy compared to triweekly therapy and concurrent treatment with gemcitabine and irinotecan. Severe pneumonitis and pulmonary fibrosis resulting in death have been described. Mild cases of pneumonitis tend to resolve spontaneously or after low-dose prednisone therapy (*ie*, prednisone, 40 mg daily for 2 weeks). Mild pneumonitis is not a contraindication to subsequent paclitaxel therapies, and the safe readministration of paclitaxel has been reported. Chest imaging findings include bilateral reticular or reticulonodular opacities, focal consolidation, and bilateral patchy areas of increased attenuation with upper lobe predominance. A hypersensitivity mechanism has been suggested in the pathogenesis on lung injury. Infusion-related reactions and hypersensitivity reactions may cause bronchospasm and hypotension.⁷²⁻⁷⁹

TOPOISOMERASE I INHIBITORS

Irinotecan

Irinotecan is a topoisomerase I inhibitor that is used mainly in the treatment of colon cancer, particularly in combination with other chemotherapy agents. Pneumonitis is a dose-dependent side effect of irinotecan. Moderate-to-severe pneumonitis has been reported in 2 to 16% of patients treated with irinotecan. Severe hypoxemia and respiratory failure requiring mechanical ventilation may develop in about 9% of the patients. Fatalities due to severe pneumonitis have been reported in 1 to 3.5% of patients.⁸⁰⁻⁸³

Topotecan

Topotecan is a topoisomerase I inhibitor that is mainly used in the treatment of metastatic carcinoma of the ovary or small cell lung cancer. Dyspnea has been reported in 3 to 4% of patients treated for ovarian or lung cancer.^{84,85} Few cases of topotecan-induced lung toxicity have been described, and include diffuse alveolar damage, organizing pneumonia, mild interstitial fibrosis with numerous intraalveolar macrophages, and respiratory failure in the setting of preexisting pulmonary fibrosis.⁸⁶⁻⁸⁸

Exatecan

Exatecan is a new and experimental topoisomerase I inhibitor that has activity against a number of solid tumors. Although in one phase II study⁸⁹ mild to moderate dyspnea developed in 36% of patients treated with exatecan, pulmonary toxicity secondary to exatecan therapy has not been reported.

THALIDOMIDE

Thalidomide was approved in 2006 for the treatment of multiple myeloma. Thalidomide is also being investigated for treatment of several other cancers. Dyspnea in association with thalidomide therapy has been reported^{90,91} in 4 to 54% of patients. Opportunistic infections including *Pneumocystis carinii* pneumonia, disseminated herpes zoster, and herpes simplex infections have been described in leukopenic patients receiving thalidomide.^{92,93} Several studies^{94,95} have found increased incidence of DVT and pulmonary embolism in association with thalidomide therapy. Thromboembolic disease usually occurs at a mean time of 2 months after thalidomide administration. The reported rates of DVT and pulmonary embolism vary from 0 to 43% of treated patients. Higher rates have been observed among patients who received thalidomide in combination with chemotherapy (16%) or dexamethasone (15%) compared to thalidomide alone (5%).^{94,95} Thalidomide-induced pneumonitis is extremely rare. Mild interstitial fibrosis, lymphocytic alveolitis, and organizing pneumonia have been described.^{4,96,97}

TYROSINE KINASE INHIBITORS

Gefitinib

Gefitinib is an oral EGFR tyrosine kinase inhibitor that is active against NSCLC, and ovarian, colon, head and neck, and breast cancers. Gefitinib-induced lung toxicity usually occurs within the first 90 days of treatment with gefitinib. Interstitial pneumonitis, diffuse alveolar damage, alveolar hemorrhage, and pulmonary fibrosis have been described.⁹⁸⁻¹⁰³ The reported incidence of gefitinib-induced lung toxicity in Japan is between 1% and 2%, which is higher than the incidence of 0.3% that has been reported in the United States.¹⁰⁰⁻¹⁰³ Gefitinib-induced pneumonitis is fatal in one third of cases.¹⁰⁴ Risk factors for pneumonitis seem to be the presence of previous lung damage from smoking, chemotherapy, irradiation, infection, or pulmonary fibrosis.²⁴

Erlotinib

Erlotinib is a HER type 1/EGFR tyrosine kinase inhibitor. Erlotinib is indicated for the treatment of patients with locally advanced or metastatic NSCLC. Erlotinib-induced pneumonitis has been described¹⁰⁵ in patients treated with erlotinib for the treatment of advanced solid tumors. In a National Cancer Institute of Canada Clinical Trials Group study,¹⁰⁶ erlotinib, 150 mg daily, was compared with placebo.

Patients with locally advanced or metastatic NSCLC after failure to respond to at least one prior chemotherapy regimen were enrolled into this study. The incidence of pneumonitis in this study was 0.8%.¹⁰⁶ In a phase III trial¹⁰⁷ of erlotinib hydrochloride combined with carboplatin and paclitaxel chemotherapy in patients with advanced NSCLC, severe pneumonitis and respiratory failure developed in five patients in the erlotinib arm (1.0%) and one patient in the placebo arm (0.2%). All six cases of pneumonitis were fatal.¹⁰⁷ The clinical presentation and nature of these interstitial lung diseases were not reported in these trials. Vahid and Esmaili¹⁰⁸ described two cases of erlotinib-induced pneumonitis that resulted in respiratory failure. The patients presented 4 to 6 days after the initiation of erlotinib therapy with fever, cough, and hypoxemia. Bilateral ground-glass opacities seen on chest imaging and elevated BAL fluid cell counts with high neutrophil percentages (76% and 89%, respectively) were found. One patient improved with high-dose corticosteroid therapy and was extubated in 4 days. Septic shock developed in the second patient, and the patient died.¹⁰⁸

Imatinib

Imatinib is a potent tyrosine kinase inhibitor that is mainly used in the treatment of patients with chronic myelogenous leukemia (CML). It is also effective in patients with GI stromal tumor. Although most cases of imatinib-induced pulmonary adverse events have been reported in patients with CML (0.2 to 1.3% in patients with early chronic phase CML), there have been rare cases of pneumonitis described in patients with GI stromal tumor treated with imatinib. Dyspnea during imatinib therapy is most often related to fluid retention and pulmonary edema. Fluid retention may be due to prolonged platelet-derived growth factor inhibition by imatinib. Platelet-derived growth factor pathways are involved in the regulation of interstitial fluid homeostasis.¹⁰⁹ Imatinib pneumonitis develops 10 to 282 days (median time, 49 days) after treatment with imatinib (range, 200 to 600 mg daily). Dyspnea, hypoxemia, fever, eosinophilia, and elevated KL-6 levels are usually seen. The chest CT scan shows diffuse or patchy ground-glass opacity, consolidation, or fine nodular opacity. The lung pathology may show interstitial pneumonitis and fibrosis, destruction of alveolar septa, lymphocytic alveolitis, plasma cell infiltrates, or type II pneumocyte hyperplasia. The resolution of pneumonitis after corticosteroid therapy has been reported. Ohnishi et al¹⁰⁹ reported that pneumonitis developed in 4 of 11 patients with a history of imatinib-induced pneumonitis after reexposure to imatinib.^{110–115} Pleural ef-

fusions (unilateral and bilateral) have been described after the initiation of imatinib therapy. It is important to acknowledge that although drug toxicity and the subsequent development of pleural effusions is possible, it is extremely difficult to exclude volume overload and the progression of primary disease as a cause of pleural effusion in reported cases. The nature of these drug-induced effusions has not been described.¹¹⁶

TREATMENT

The mainstay of therapy for drug-induced pneumonitis is the cessation of the presumed culprit agent and systemic corticosteroids. Although corticosteroids are used widely to treat drug-induced pneumonitis, this treatment has not been evaluated in controlled clinical trials. It is important to exclude an infectious etiology prior to initiating corticosteroid therapy. We recommend therapy with methylprednisolone, 1 g/d for 3 days, in patients with respiratory failure. Lower doses of corticosteroids (methylprednisolone, 60 mg every 6 h) may be used in less severe cases of pneumonitis.^{1,2} In our experience, therapy with systemic corticosteroids results in a rapid improvement in oxygenation and may lead to mechanical ventilation liberation. Supportive care, bronchodilators, IV fluid, vasopressors, and mechanical ventilation are indicated in patients with severe hypersensitivity reactions and circulatory collapse.

REFERENCES

- 1 Higenbottam T, Kuwano K, Nemery B, et al. Understanding the mechanism of drug-associated interstitial lung disease. *Br J Cancer* 2004; 91(suppl):S31–S37
- 2 Camus P, Kudoh S, Ebina M. Interstitial lung disease associated with drug therapy. *Br J Cancer* 2004; 91(suppl): S18–S23
- 3 Michielin O, Udry E, Periard D, et al. Irinotecan-induced interstitial pneumonia. *Lancet Oncol* 2004; 5:322–324
- 4 Feaver AA, McCune DE, Mysliwiec AG, et al. Thalidomide-induced organizing pneumonia. *South Med J* 2006; 99:1292–1294
- 5 Athanasiadis A, Roussos G, Papakostoulis T, et al. Paclitaxel and gemcitabine combination in a biweekly schedule in patients with advanced non small-cell lung cancer: a phase I study. *Cancer Chemother Pharmacol* 2005; 56:653–658
- 6 Okuno SH, Frytak S. Mitomycin lung toxicity: acute and chronic phase. *Am J Clin Oncol* 1997; 20:282–284
- 7 Schrijvers D, Catimel G, Highley M, et al. KW-2149-induced lung toxicity is not prevented by corticosteroids: a phase I and pharmacokinetic study. *Anticancer Drugs* 1999; 10:633–639
- 8 Kobayashi J, Kitamura S. KL-6: a serum marker for interstitial pneumonia. *Chest* 1995; 108:311–315
- 9 Ohnishi H, Yokoyama A, Yasuhara Y, et al. Circulating KL-6 levels in patients with drug induced pneumonitis. *Thorax* 2003; 58:872–875

- 10 Kobayashi J, Kitamura S. Serum KL-6 for the evaluation of active pneumonitis in pulmonary sarcoidosis. *Chest* 1996; 109:1276–1282
- 11 Kase S, Kitaichi N, Namba K, et al. Elevation of serum Krebs von den Lunge-6 levels in patients with tubulointerstitial nephritis and uveitis syndrome. *Am J Kidney Dis* 2006; 48:935–941
- 12 Sato H, Callister MEJ, Mumby S, et al. KL-6 levels are elevated in plasma from patients with acute respiratory distress syndrome. *Eur Respir J* 2004; 23:142–145
- 13 Ogawa Y, Ishikawa T, Ikeda K, et al. Evaluation of serum KL-6, a mucin-like glycoprotein, as a tumor marker for breast cancer. *Clin Cancer Res* 2000; 6:4069–4072
- 14 Kawata T, Tsukagoshi H, Mashimo T, et al. KL-6-producing invasive thymoma. *Intern Med* 2002; 41:979–982
- 15 Kurosaki M, Izumi N, Onuki Y, et al. Serum KL-6 as a novel tumor marker for hepatocellular carcinoma in hepatitis C virus infected patients. *Hepatol Res* 2005; 33:250–257
- 16 Miya T, Ono Y, Tanaka H, et al. Radiation recall pneumonitis induced by gefitinib (Iressa): a case report. *Nihon Kokyuki Gakkai Zasshi* 2003; 41:565–568
- 17 Thomas PS, Agrawal S, Gore M, et al. Recall lung pneumonitis due to carmustine after radiotherapy. *Thorax* 1995; 50:1116–1118
- 18 Schweitzer VG, Juillard GJ, Bajada CL, et al. Radiation recall dermatitis and pneumonitis in a patient treated with paclitaxel. *Cancer* 1995; 76:1069–1072
- 19 Ma LD, Taylor GA, Wharam MD, et al. “Recall” pneumonitis: adriamycin potentiation of radiation pneumonitis in two children. *Radiology* 1993; 187:465–467
- 20 Kataoka M, Kawamura M, Nishiyama Y, et al. A case with delayed-onset radiation pneumonitis suspected to be induced by oral etoposide. *Nippon Igaku Hoshasen Gakkai Zasshi* 1992; 52:641–645
- 21 Vegesna V, Withers HR, McBride WH, et al. Adriamycin-induced recall of radiation pneumonitis and epilation in lung and hair follicles of mouse. *Int J Radiat Oncol Biol Phys* 1992; 23:977–981
- 22 Hill AB, Tattersall SF. Recall of radiation pneumonitis after intrapleural administration of doxorubicin. *Med J Aust* 1983; 1:39–40
- 23 Barlési F, Villani P, Doddoli C, et al. Gemcitabine-induced severe pulmonary toxicity. *Fundam Clin Pharmacol* 2004; 18:85–91
- 24 Takano T, Ohe Y, Kusumoto M, et al. Risk factors for interstitial lung disease and predictive factors for tumor response in patients with advanced non-small cell lung cancer treated with gefitinib. *Lung Cancer* 2004; 45:93–104
- 25 Atabai K, Ishigaki M, Geiser T, et al. Keratinocyte growth factor can enhance alveolar epithelial repair by nonmitogenic mechanisms. *Am J Physiol* 2002; 283:163–169
- 26 Miettinen PJ, Warburton D, Bu D, et al. Impaired lung branching morphogenesis in the absence of functional EGF receptor. *Dev Biol* 1997; 186:224–236
- 27 Dimopoulou I, Bamias A, Lyberopoulos P, et al. Pulmonary toxicity from novel antineoplastic agents. *Ann Oncol* 2006; 17:372–379
- 28 Sordillo EM, Sordillo PP, Stover D, et al. Chlorozotocin (DCNU)-induced pulmonary toxicity. *Cancer Clin Trials* 1981; 4:397–399
- 29 Geodert JJ, Smith FP, Tsou E, et al. Combination chemotherapy pneumonitis: a case report of possible synergistic toxicity. *Pediatr Oncol* 1983; 11:116–118
- 30 Baker WJ, Fistel SJ, Jones RV, et al. Interstitial pneumonitis associated with ifosfamide therapy. *Cancer* 1990; 65:2217–2221
- 31 Chen YM, Shih JF, Lee CS, et al. Phase II study of docetaxel and ifosfamide combination chemotherapy in non-small-cell lung cancer patients failing previous chemotherapy with or without paclitaxel. *Lung Cancer* 2003; 39:209–214
- 32 Hadjiladis D, Govert JA. Methemoglobinemia after infusion of ifosfamide chemotherapy. *Chest* 2000; 118:1208–1210
- 33 Pasetto LM, Monfardini S. Is acute dyspnea related to oxaliplatin administration? *World J Gastroenterol* 2006; 12:5907–5908
- 34 Yague XH, Soy E, Merino BQ, et al. Interstitial pneumonitis after oxaliplatin treatment in colorectal cancer. *Clin Transl Oncol* 2005; 7:515–517
- 35 Gagnadoux F, Roiron C, Carrie E, et al. Eosinophilic lung disease under chemotherapy with oxaliplatin for colorectal cancer. *Am J Clin Oncol* 2002; 25:388–390
- 36 Lee MY, Yang MH, Liu JH, et al. Severe anaphylactic reactions in patients receiving oxaliplatin therapy: a rare but potentially fatal complication. *Support Care Cancer* 2007; 15:89–93
- 37 Abrey LE, Oslon JD, Raizer JJ, et al. A phase II trial of temozolomide for patients with recurrent or progressive brain metastasis. *J Neurooncol* 2001; 53:259–265
- 38 Brandwein JM, Yang L, Schimmer AD, et al. A phase II study of temozolomide therapy for poor-risk patients aged ≥ 60 years with acute myeloid leukemia: low levels of MGMT predict for response. *Leukemia* 2007; 21:821–824
- 39 Skubitz KM, Skubitz APN. Mechanism of transient dyspnea induced by pegylated-liposomal doxorubicin (Doxil). *Anti Cancer Drugs* 1998; 9:45–50
- 40 Tsao YT, Dai MS, Chang H, et al. Bronchiolitis obliterans organizing pneumonia presenting as hemoptysis in a patient of Hodgkin’s lymphoma undergoing chemotherapy. *J Med Sci* 2006; 26:115–118
- 41 Jacobs C, Slade M, Lavery B. Doxorubicin and BOOP: a possible near fatal association. *Clin Oncol* 2002; 14:262
- 42 Dang CT, D’Andrea G, Moynahan ME, et al. Phase II study of feasibility of dose-dense FEC followed by alternating weekly taxanes in high-risk, four or more node-positive breast cancer. *Clin Cancer Res* 2004; 10:5754–5761
- 43 Tomlinson J, Tighe M, Johnson S, et al. Interstitial pneumonitis following mitoxantrone, chlorambucil and prednisolone (MCP) chemotherapy. *Clin Oncol* 1999; 11:184–186
- 44 Quigley M, Brada M, Heron C, et al. Severe lung toxicity with a weekly low dose chemotherapy regimen in patients with non-Hodgkin’s lymphoma. *Hematol Oncol* 1988; 6: 319–324
- 45 Matsukawa Y, Takeuchi J, Aiso M, et al. Interstitial pneumonitis possibly due to mitoxantrone. *Acta Haematol* 1993; 90:155–158
- 46 Roth BJ, Manola J, Dreicer R, et al. Piritrexim in advanced, refractory carcinoma of the urothelium (E3896): a phase II trial of the Eastern Cooperative Oncology Group. *Invest New Drugs* 2002; 20:425–429
- 47 de Wit R, Verweij J, Slingerland R, et al. Piritrexim-induced pulmonary toxicity. *Am J Clin Oncol* 1993; 16:146–148
- 48 Johnson DH, Fehrenbacher L, Novotny WF, et al. Randomized phase II trial comparing bevacizumab plus carboplatin and paclitaxel with carboplatin and paclitaxel alone in previously untreated locally advanced or metastatic non-small-cell lung cancer. *J Clin Oncol* 2004; 22:2184–2191
- 49 Sandler A, Gray R, Perry MC, et al. Paclitaxel-carboplatin alone or with bevacizumab for non-small-cell lung cancer. *N Engl J Med* 2006; 355:2542–2550
- 50 Herbst RS, Sandler AB. Non-small cell lung cancer and antiangiogenic therapy: what can be expected of bevacizumab? *Oncologist* 2004; 9(suppl):19–26

- 51 Kollmannsberger C, Schttenhelm M, Honecker F, et al. A phase I study of the humanized monoclonal anti-epidermal growth factor receptor (EGFR) antibody EMD 72000 (Matuzumab) in combination with paclitaxel in patients with EGFR-positive advanced non-small-cell lung cancer (NSCLC). *Ann Oncol* 2006; 17:1007–1013
- 52 Vahid B, Mehrotra A. Trastuzumab (herceptin)-associated lung injury. *Respirology* 2006; 11:655–658
- 53 Radzikowska E, Szczepulska E, Chabowski M, et al. Organising pneumonia caused by trastuzumab (herceptin) therapy for breast cancer. *Eur Respir J* 2003; 21:552–555
- 54 Tripathy D, Slamon DJ, Cobleigh M, et al. Safety of treatment of metastatic breast cancer with trastuzumab beyond disease progression. *J Clin Oncol* 2004; 22:1063–1070
- 55 Clamon G, Herndon J, Kern J, et al. Lack of trastuzumab activity in nonsmall cell lung carcinoma with overexpression of erb-B2:39810: a phase II trial of Cancer and Leukemia Group B. *Cancer* 2005; 103:1670–1675
- 56 Romond EH, Perez EA, Bryant J, et al. Trastuzumab plus adjuvant chemotherapy for operable HER2-positive breast cancer. *N Engl J Med* 2005; 353:1673–1684
- 57 Briasoulis E, Pavlidis. Noncardiogenic pulmonary edema: an unusual and serious complication of anticancer therapy. *Oncologist* 2001; 6:153–161
- 58 Roychowdhury DF, Cassidy CA, Peterson P, et al. A report on serious pulmonary toxicity associated with gemcitabine-based therapy. *Invest New Drugs* 2002; 20:311–315
- 59 Gupta N, Ahmed I, Steinberg H, et al. Gemcitabine-induced pulmonary toxicity. *Am J Clin Oncol* 2002; 25:96–100
- 60 Marruchella A, Fiorenzano G, Merizzi A, et al. Diffuse alveolar damage in a patient treated with gemcitabine. *Eur Respir J* 1998; 11:504–506
- 61 Pavlaski N, Bell DR, Millward MJ, et al. Fatal pulmonary toxicity resulting from treatment with gemcitabine. *Cancer* 1997; 80:286–291
- 62 Carron P-L, Cousin L, Caps T, et al. Gemcitabine-associated diffuse alveolar damage. *Intensive Care Med* 2001; 27:1554
- 63 Vander Els NJ, Miller V. Successful treatment of gemcitabine toxicity with a brief course of oral corticosteroid therapy. *Chest* 1998; 114:1779–1781
- 64 Siderov J, Prasad P, De Boer R, et al. Safe administration of etoposide phosphate after hypersensitivity reaction to intravenous etoposide. *Br J Cancer* 2002; 86:12–13
- 65 Dajczman E, Srolovitz H, Kreisman H, et al. Fatal pulmonary toxicity following oral etoposide therapy. *Lung Cancer* 1995; 12:81–86
- 66 Gurjal A, An T, Valdivieso M, et al. Etoposide-induced pulmonary toxicity. *Lung Cancer* 1999; 26:109–112
- 67 Zimmerman MS, Ruckdeschel JC, Hussain M. Chemotherapy-induced interstitial pneumonitis during treatment of small cell anaplastic lung cancer. *J Clin Oncol* 1984; 2:396–405
- 68 O'Dwyer PJ, King SA, Fortner CL, et al. Hypersensitivity reactions to teniposide (VM-26): an analysis. *J Clin Oncol* 1986; 8:1262–1269
- 69 Duran I, Siu LL, Oza AM, et al. Characterization of the lung toxicity of the cell cycle inhibitor temsirolimus. *Eur J Cancer* 2006; 42:1875–1880
- 70 Atkins MB, Hidalgo M, Stadler WM, et al. Randomized phase II study of multiple dose levels of CCI-779, a novel mammalian target of rapamycin kinase inhibitor, in patients with advanced refractory renal cell carcinoma. *J Clin Oncol* 2004; 22:909–918
- 71 Rothenburger M, Teerling E, Bruch C, et al. Calcineurin inhibitor-free immunosuppression using everolimus (certican) in maintenance heart transplant recipients: 6 months' follow-up. *J Heart Lung Transplant* 2007; 26:250–257
- 72 Shitara K, Ishii E, Kondo M, et al. Suspected paclitaxel-induced pneumonitis. *Gastric Cancer* 2006; 9:325–328
- 73 Suzuki N, Hiraki A, Takigawa N, et al. Severe interstitial pneumonia induced by paclitaxel in a patient with adenocarcinoma of the lung. *Acta Med Okayama* 2006; 60:295–298
- 74 Ostoros G, Pretz A, Fillinger J, et al. Fatal pulmonary fibrosis induced by paclitaxel: a case report and review of the literature. *Int J Gynecol Cancer* 2006; 16(suppl):391–393
- 75 Takahashi T, Higashi S, Nishiyama H, et al. Biweekly paclitaxel and gemcitabine for patients with advanced urothelial cancer ineligible for cisplatin-based regimen. *Jpn J Clin Oncol* 2006; 36:104–108
- 76 Leimgruber K, Negro R, Baier S, et al. Fatal interstitial pneumonitis associated with docetaxel administration in a patient with hormone-refractory prostate cancer. *Tumori* 2006; 92:542–544
- 77 Kouroussis C, Mavroudis D, Kakolyris S, et al. High incidence of pulmonary toxicity of weekly docetaxel and gemcitabine in patients with non-small cell lung cancer: results of a dose-finding study. *Lung Cancer* 2004; 44:363–368
- 78 Ramanathan R, Reddy VV, Holbert JM, et al. Pulmonary infiltrates following administration of paclitaxel. *Chest* 1996; 110:289–292
- 79 Wong P, Leung AN, Berry GJ, et al. Paclitaxel-induced hypersensitivity pneumonitis: radiographic and CT findings. *Am J Roentgenol* 2001; 176:718–720
- 80 Rocha-Lima CM, Herndon JE II, Lee ME, et al. Phase II trial of irinotecan/gemcitabine as second-line therapy for relapsed and refractory small-cell lung cancer: Cancer and Leukemia Group B Study 39902. *Ann Oncol* 2007; 18:331–337
- 81 Kaneda H, Kurata T, Tamura K, et al. A phase I study of irinotecan in combination with amrubicin for advanced lung cancer patients. *Anticancer Res* 2006; 26:2479–2485
- 82 Klautke G, Fahndrich S, Semrau S, et al. Simultaneous chemoradiotherapy with irinotecan and cisplatin in limited disease small cell lung cancer: a phase I study. *Lung Cancer* 2006; 53:183–188
- 83 Madarnas Y, Webster P, Shorter AM, et al. Irinotecan-associated pulmonary toxicity. *Anticancer Drugs* 2000; 11:709–713
- 84 O'Brien ME, Ciuleanu TE, Tsekov H, et al. Phase III trial comparing supportive care alone with supportive care with oral topotecan in patients with relapsed small-cell lung cancer. *J Clin Oncol* 2006; 24:5441–5447
- 85 Pfisterer J, Weber B, Reuss A, et al. Randomized phase III trial of topotecan following carboplatin and paclitaxel in first-line treatment of advanced ovarian cancer: a Gynecologic Cancer Intergroup Trial of the AGO-OVAR and GINECO. *J Natl Cancer Inst* 2006; 98:1036–1045
- 86 Edgerton CC, Gilman M, Roth BJ. Topotecan-induced bronchiolitis. *South Med J* 2007; 97:699–701
- 87 Rossi SE, Erasmus JJ, McAdams HP, et al. Pulmonary drug toxicity: radiologic and pathologic manifestations. *Radiographics* 2000; 20:1245–1259
- 88 Maitland ML, Wilcox R, Hogarth DK, et al. Diffuse alveolar damage after a single dose of topotecan in a patient with pulmonary fibrosis and small cell cancer. *Lung Cancer* 2006; 54:243–245
- 89 Reichardt P, Nielsen OS, Bauer S, et al. Exatecan in pretreated adult patients with advanced soft tissue sarcoma: results of a phase II-study of the EORTC Soft Tissue and Bone Sarcoma Group. *Eur J Cancer* 2007; 43:1017–1022
- 90 Gordinier ME, Dizon DS. Dyspnea during thalidomide

- treatment for advanced ovarian cancer. *Ann Pharmacother* 2005; 39:962–965
- 91 Rajkumar SV, Hayman S, Gertz MA, et al. Combination therapy with thalidomide plus dexamethasone for newly diagnosed myeloma. *J Clin Oncol* 2002; 20:4319–4323
 - 92 Curley MJ, Hussein SA, Hassoun PM. Disseminated herpes simplex virus and varicella zoster virus coinfection in a patient taking thalidomide for relapsed multiple myeloma. *J Clin Microbiol* 2002; 40:2302–2304
 - 93 Kulke MH, Stuart K, Enzinger PC, et al. Phase II study of temozolomide and thalidomide in patients with metastatic neuroendocrine tumors. *J Clin Oncol* 2006; 24:401–406
 - 94 Kumar S, Witzig TE, Rajkumar SV. Thalidomide: current role in the treatment of non-plasma cell malignancies. *J Clin Oncol* 2004; 22:2477–2488
 - 95 Bennett CL, Schumock GT, Desai AA, et al. Thalidomide-associated deep venous thrombosis and pulmonary embolism. *Am J Med* 2002; 113:603–606
 - 96 Iguchi T, Sakoda M, Chen CK, et al. Interstitial pneumonia during treatment with thalidomide in a patient with multiple myeloma. *Rinsho Ketsueki* 2004; 45:1064–1066
 - 97 Onozawa M, Hashino S, Sogabe S, et al. Side effects and good effects from new chemotherapeutic agents: case 2. Thalidomide-induced interstitial pneumonitis. *J Clin Oncol* 2005; 23:2425–2426
 - 98 Inoue A, Saijo Y, Maemondo M, et al. Severe acute interstitial pneumonia and gefitinib. *Lancet* 2003; 361:137–139
 - 99 Okamoto I, Fujii K, Matsumoto M, et al. Diffuse alveolar damage after ZD 1839 therapy in a patient with non-small cell lung cancer. *Lung Cancer* 2003; 40:339–342
 - 100 Ohyanagi F, Ando Y, Nagashima F, et al. Acute gefitinib-induced pneumonitis. *Int J Clin Oncol* 2004; 9:406–409
 - 101 Sumpter K, Harper-Wynne C, O'Brien M, et al. Severe acute interstitial pneumonia and gefitinib. *Lung Cancer* 2004; 43:367–368
 - 102 Ieki R, Saitoh, Shibuya M. Acute lung injury as a possible adverse drug reaction related to gefitinib. *Eur Respir J* 2003; 22:179–181
 - 103 Nagaria NC, Cogswell J, Choe JK, et al. Side effects and good effects from new chemotherapeutic agents: case 1. Gefitinib-induced interstitial fibrosis. *J Clin Oncol* 2005; 23:2423–2424
 - 104 Cohen MH, Williams GA, Sridhara R, et al. FDA drug approval summary: gefitinib (ZD1839) (Iressa) tablets. *Oncologist* 2003; 8:303–306
 - 105 Tammaro KA, Baldwin PD, Lundberg AS. Interstitial lung disease following erlotinib (Tarceva) in a patient who previously tolerated gefitinib (Iressa). *J Oncol Pharm Pract* 2005; 11:127–130
 - 106 Herbst RS, Prager D, Hermann R, et al. TRIBUTE: a phase III trial of erlotinib hydrochloride (OSI-774) combined with carboplatin and paclitaxel chemotherapy in advanced non-small-cell lung cancer. *J Clin Oncol* 2005; 23:5892–5899
 - 107 Shepherd FA, Rodrigues Pereira J, Ciuleanu T, et al. Erlotinib in previously treated non-small-cell lung cancer. *N Engl J Med* 2005; 353:123–132
 - 108 Vahid B, Esmaili A. Erlotinib-associated acute pneumonitis: report of two cases. *Can Respir J* 2007; 14:167–170
 - 109 Ohmishi K, Sakai F, Kudoh S, et al. Twenty-seven cases of drug-induced interstitial lung disease associated with imatinib. *Leukemia* 2006; 20:1162–1164
 - 110 Lin JT, Yeh KT, Fang HY, et al. Fulminant, but reversible interstitial pneumonitis associated with imatinib mesylate. *Leuk Lymphoma* 2006; 47:1693–1695
 - 111 Rajda J, Phatak PD. Reversible drug-induced interstitial pneumonitis following imatinib mesylate therapy. *Am J Hematol* 2005; 79:80–81
 - 112 Grimison P, Goldstein D, Schneeweiss J, et al. Corticosteroid-responsive interstitial pneumonitis related to imatinib mesylate with successful rechallenge, and potential causative mechanisms. *Intern Med J* 2005; 35:136–137
 - 113 Isshiki I, Yamaguchi K, Okamoto S. Interstitial pneumonitis during imatinib therapy. *Br J Haematol* 2004; 125:420
 - 114 Ma CX, Hobday TJ, Jett JR. Imatinib mesylate-induced interstitial pneumonitis. *Mayo Clin Proc* 2003; 78:1578–1579
 - 115 Bergeron A, Bergot E, Vilela G, et al. Hypersensitivity pneumonitis related to imatinib mesylate. *J Clin Oncol* 2002; 20:4271–4272
 - 116 Goldsby R, Pulsipher M, Adams R, et al. Unexpected pleural effusions in 3 pediatric patients treated with STI-571. *J Pediatr Hematol Oncol* 2002; 24:694–695

Pulmonary Complications of Novel Antineoplastic Agents for Solid Tumors

Bobbak Vahid and Paul E. Marik
Chest 2008;133:528-538
DOI 10.1378/chest.07-0851

This information is current as of July 10, 2008

Updated Information & Services	Updated information and services, including high-resolution figures, can be found at: http://chestjournal.org/cgi/content/full/133/2/528
References	This article cites 116 articles, 41 of which you can access for free at: http://chestjournal.org/cgi/content/full/133/2/528#BIBL
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://chestjournal.org/misc/reprints.shtml
Reprints	Information about ordering reprints can be found online: http://chestjournal.org/misc/reprints.shtml
Email alerting service	Receive free email alerts when new articles cite this article sign up in the box at the top right corner of the online article.
Images in PowerPoint format	Figures that appear in CHEST articles can be downloaded for teaching purposes in PowerPoint slide format. See any online article figure for directions.

A M E R I C A N C O L L E G E O F



P H Y S I C I A N S[®]